



# Drug Therapy Protocols: Hydroxocobalamin

<b>Policy code</b>	DTP_HYDX_o626
<b>Date</b>	June, 2026
<b>Purpose</b>	To ensure a consistent procedural approach to hydroxocobalamin administration.
<b>Scope</b>	Applies to all Queensland Ambulance Service (QAS) clinical staff.
<b>Health care setting</b>	Pre-hospital assessment and treatment.
<b>Population</b>	Applies to all ages unless specifically mentioned.
<b>Source of funding</b>	Internal – 100%
<b>Author</b>	Clinical Quality & Patient Safety Unit, QAS
<b>Review date</b>	June, 2028
<b>Information security</b>	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
<b>URL</b>	<a href="https://ambulance.qld.gov.au/clinical.html">https://ambulance.qld.gov.au/clinical.html</a>

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.


All feedback and suggestions are welcome. Please forward to: [Clinical.Guidelines@ambulance.qld.gov.au](mailto:Clinical.Guidelines@ambulance.qld.gov.au)

## Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by appropriately qualified QAS clinicians when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2026.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: [Clinical.Guidelines@ambulance.qld.gov.au](mailto:Clinical.Guidelines@ambulance.qld.gov.au)

# Hydroxocobalamin

July, 2022

## Drug class

Antidote<sup>[1,2]</sup>

## Pharmacology

Hydroxocobalamin (an injectable form of vitamin B12) is an antidote for cyanide toxicity. It binds with circulating and cellular cyanide to form cyanocobalamin, which is then excreted in urine.<sup>[1,2]</sup>

## Metabolism

Excreted by the kidneys<sup>[1]</sup>

### Indications

- **Life-threatening cyanide toxicity** (e.g. shock, respiratory failure, seizure, ALOC or myocardial ischaemia)

### Contraindications

- Allergy AND/OR Adverse Drug Reaction

### Precautions

- Hypertension (*refer to Special notes*)
- Pregnancy (*refer to Special notes*)

### Side effects

- Anaphylaxis
- Chromaturia
- Erythema
- Rash (acne like)
- Hypertension
- Renal failure
- Headache
- Nausea and/or vomiting
- Pain at infusion site

### Presentation

- Vial, 5 g hydroxocobalamin (CYANOKIT®)<sup>[3]</sup>

### Onset (IV)

Immediate

### Duration (IV)

Several days

### Half-life

26–31 hours

## Schedule

- N/A – TGA Special Access Scheme.

## Routes of administration

Intravenous infusion (IV INF)



## Special notes

- Hydroxocobalamin infusions are only to be administered by appropriately trained QAS clinicians within the following response catchments:
  - *Central Queensland District:* Orica Yarwun Cyanide Plant
  - *Darling Downs District:* Texas Silver Mine
  - *North West District:* Great Australian Mine, Lorena Gold Project, Mt Isa Mines, George Fisher
- For patients presenting with non-life threatening toxicity (e.g. headache, confusion, vomiting, dyspnoea or chest tightness) the *QAS Clinical Consultation & Advice Line* should be immediately contacted regarding potential hydroxocobalamin administration.
- Substantial increases in blood pressure may occur following hydroxocobalamin therapy.
- There are no adequate and well-controlled studies of hydroxocobalamin administration in pregnant women. Hydroxocobalamin should be used during pregnancy **ONLY** if the potential benefits justifies the potential risk to the fetus.

## Special notes (cont.)

- Each CYANOKIT® contains the following components:
  - one 250 mL glass vial containing 5 g lyophilised hydroxocobalamin for injection,
  - one sterile transfer spike,
  - one sterile vented infusion set,
  - one quick use reference guide and one package insert.
- **\* NOTE: dilutant is NOT included (2 x 100 mL sodium chloride 0.9% required).**
- Simultaneous administration of hydroxocobalamin with other medications and/or blood products is not recommended.
- All parenteral medications must be prepared in an aseptic manner. The rubber stopper of all vials must be disinfected with an appropriate antimicrobial swab and allowed to dry prior to piercing.
- All hydroxocobalamin infusions must be initiated using industry supplied stock. Hydroxocobalamin will not be procured by QAS.
- All cannulae and IV lines must be flushed thoroughly with sodium chloride 0.9% following each medication administration.

## Adult dosages<sup>[1-4]</sup>

**Life-threatening cyanide toxicity** (e.g. shock, respiratory failure, seizure, ALOC or myocardial ischaemia)



**IV  
INF**


**5 g over 15 minutes.**

## Paediatric dosages

**Note:** QAS officers are **NOT** authorised to administer hydroxocobalamin to paediatric patients.

## Hydroxocobalamin (CYANOKIT®) preparation


1. Keeping the CYANOKIT® 5g vial in the supplied packaging, place it on a firm flat surface with the cap uppermost. Remove the cap and disinfect the rubber stopper with a 2% chlorhexidine / 70% isopropyl alcohol swab and allow to dry.



2. Take the supplied transfer spike from the packaging and remove the safety cap from one end.



3. Remove the safety cap from one of the 100 mL sodium chloride 0.9% bags and firmly insert the exposed transfer spike.
4. Remove the remaining safety cap from the transfer spike and while maintaining aseptic technique, hold the transfer spike's wings, allow the bag to hang down (so as to not spill any contents) and insert the exposed transfer spike into the rubber stopper of the upright CYANOKIT® vial.




5. Lift the sodium chloride 0.9% bag to empty it fully into the CYANOKIT® vial.



6. While leaving the transfer spike in the upright CYANOKIT® vial gently remove the empty sodium chloride bag.



7. Remove the safety cap from the second 100 mL sodium chloride 0.9% bag and while maintaining aseptic technique carefully insert the exposed transfer spike into the second sodium chloride bag, keeping the CYANOKIT® vial in the upright position.



8. Lift the sodium chloride 0.9% bag to empty it fully into the CYANOKIT® vial.
9. Gently remove the transfer spike with the attached sodium chloride bag from the CYANOKIT® vial.



10. Carefully mix the CYANOKIT® vial solution by repeatedly inverting or rocking the vial for 60 seconds; **do not shake the vial.**
11. Rotate the CYANOKIT® vial within the carton until the glass is viewable in the window. Inspect to confirm the solution is cherry red and free of particulates. Do not administer if it is NOT cherry red or if it contains particulates (contact the QAS Clinical Consultation & Advice Line for advice).



12. Prime a QAS supplied Alaris™ (gravity flow) giving set with the reconstituted medication and ensure no air bubbles are present in the line. Open the giving set's red air vent and administer in accordance with DTP: Hydroxocobalamin.