



Clinical Practice Guidelines: Medical/Diabetic emergency: Hypoglycaemia

Policy code	CPG_ME_DHO_0326
Date	March, 2026
Purpose	To ensure consistent management of patients with hypoglycaemia.
Scope	Applies to Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless stated otherwise.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	March, 2029
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by appropriately qualified QAS clinicians when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2026.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Diabetic emergency: Hypoglycaemia

March, 2026

Glucose is an essential metabolic fuel for the brain and a constant supply is critical for normal neurological function. **Hypoglycaemia** is defined as a BGL < 4.0 mmol/L and this can occur in any patient, regardless of a history of diabetes.^[1]

Intravenous glucose is the recommended first line management strategy in patients unable to swallow oral glucose and treatment should aim to achieve a BGL of 4.0–8 mmol/L. If there is no improvement in conscious state following such an increase in BGL, other causes for the ALOC should be considered.

Clinical features



Autonomic features (*warning signs*)

- Diaphoresis, hunger, tingling around the mouth, tremor, tachycardia, pallor, palpitations and anxiety.
- These warning signs may be lost in patients with repeated or prolonged hypoglycaemia.^[2]

Neurological features

- *Consider hypoglycaemia in all patients who have an ALOC.*
- Lethargy, change in behaviour, headache, visual disturbance, slurred speech, dizziness, ALOC, seizures, coma.
- Patients may present with signs/symptoms mimicking intoxication or stroke.

Clinical features (*cont.*)



Other considerations

- Chronic, poorly controlled diabetics may be relatively hypoglycaemic despite having a BGL greater than 4.0 mmol/L.^[2]
- Signs of hypoglycaemia may be masked in patients taking beta blocker medications.^[3]

Risk assessment



- Caution is required if the patient is agitated, aggressive or violent.
- Consideration should be given to the possibility of an accidental, or intentional hypoglycaemic agent medication overdose.

Additional information

- Patients < 2 years of age and symptomatically hypoglycaemic may be administered buccal glucose gel. See *CPP: Buccal* for further information.

Diabetes service referral

- Diabetes Service Referral must be considered for all patients (irrespective of whether transported or not) who present with diabetic related complications, e.g. hypo/hyperglycaemia.

Mandatory transport criteria

If the patient has ANY of the following, they must be transported to hospital.

- Newly diagnosed diabetes
- No previous diagnosis of diabetes
- Pregnant
- Patient recovered but unable to be monitored by a responsible adult for 4 hours, or patient unable to self-care
- Not returned to normal mental state within 10 minutes of IV glucose, or incomplete recovery to normal conscious state
- Intentional overdose of glucose lowering agent
- Severe hypoglycaemia episode within previous 48 hours
- Risk of prolonged or recurrent hypoglycaemia
 - Unwitnessed onset or prolonged episode
 - Patient taking long acting oral hypoglycaemic agent that causes hypoglycaemia
 - Unable or unwilling to consume long acting carbohydrate
- Seizure
- Suspected cause of hypoglycaemia due to illness that requires further investigation
- Injury sustained from hypoglycaemic episode requiring further investigation

Fitness to drive considerations

In alignment with the *National Driver Medication Standards*^[4], patients that experience a severe hypoglycaemic event (defined as an episode requiring intervention from a bystander or paramedic to resolve) should be advised to not operate a motor vehicle until they have been cleared to drive by a general practitioner or endocrinologist. In instances where the patient is provided this advice and attempts to drive, the Queensland Police Service should be notified. All instances where advice not to drive has been provided to patient's must be clearly documented on the eARF.

Hypoglycaemic patients with in-situ insulin pumps

In instances that patients present with a hypoglycaemic episode and have an indwelling insulin pump in-situ, ambulance clinicians are advised not to manually cease or alter the settings in these devices. However, if the patient or their carer are familiar with the workings of the pump and its settings, they may themselves decide to cease or alter the pump settings if part of a personal diabetes management plan.

The receiving facility must be advised of the patient's in-situ insulin pump on hospital arrival, along with any problems with the device that are observed by clinicians or reported by the patient or carer. This patient management recommendation has been made following a careful consideration or a risk-benefit analysis, together with external advice provided by the Queensland Health Diabetes Clinical Network.

UNCONTROLLED WHEN PRINTED

UNCONTROLLED WHEN PRINTED

UNCONTROLLED WHEN PRINTED

UNCONTROLLED WHEN PRINTED

UNCONTROLLED WHEN PRINTED

Note: Clinicians must only perform procedures for which they have received specific training and authorisation by the QAS.

* Advise patient to consume complex carbohydrates (e.g. sandwich) following oral glucose.

CPG: Clinician safety
CPG: Standard cares

Patient able to tolerate oral medication?

Consider:
• Glucose gel* (PO/BUCC)

Vascular access obtained?

Consider:
• Glucagon

Consider:
• Glucose 10%^[5,6]

If ANY of the mandatory transport criteria apply and the patient is not being transported, contact the QAS Clinical Consultation & Advice Line for case specific management.

Provide advice not to drive until medically cleared

Consider:
• Diabetes Service Referral

Determine Disposition Pathway
Pre-notify as appropriate