



Drug Therapy Protocols: Lidocaine gel 2%

Policy code	DTP_LID2_0626
Date	June, 2026
Purpose	To ensure a consistent procedural approach to lidocaine gel 2% administration.
Scope	Applies to all Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless specifically mentioned.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	June, 2028
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by appropriately qualified QAS clinicians when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2026.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Lidocaine 2% gel

March, 2026

Drug class

Local anaesthetic.

Pharmacology

Lidocaine is a local anaesthetic which produces a reversible loss of sensation by preventing or diminishing the conduction of sensory nerve impulses near the site of application.

Metabolism

Primarily metabolised by the liver and the remainder is excreted by the kidneys.

Indications



- To **provide local anaesthesia and lubrication** during urinary catheter insertion.

Contraindications



- Allergy AND/OR Adverse Drug Reaction
- Inflammation or infection of the site

Precautions



- Nil

Side effects



- Local site reactions (itch, burning sensation, rash)

Presentation

- Pre-filled syringe, 11 mL *lidocaine hydrochloride 2% gel* (InstillaGel® Lido)

Onset

3–5 minutes

Duration

20–30 minutes

Half-life

Not applicable

Schedule

- S2

Routes of administration

Topical (urethral)



Special notes

- Ambulance officers must only administer medications for the listed indications and dosing range. Any consideration for treatment outside the listed scope of practice requires mandatory approval via the *QAS Clinical Consultation and Advice Line*.
- Lidocaine 2% is to be applied for the purposes of indwelling catheter (IDC) management. For more detail, refer to *CPP: Other/Indwelling urinary catheter insertion*.
- InstillaGel® Lido gel contains a sterile lubricant.
- Males may require 20 mL (i.e., more than 1 pre-filled syringe) for adequate analgesia, the patient's age and physical status should be considered in calculating the required dose.

Adult dosages

To **provide local anaesthesia and lubrication** during urinary catheter insertion



**Topical
(urethral)**

Females: 1 pre-filled syringe (11 mL)

Apply **3–5 minutes** before insertion of catheter.

Males: 1–2 pre-filled syringe(s) (11 mL–22 mL)

Apply **3–5 minutes** before insertion of catheter.

Paediatric dosages

Note: QAS officers are **NOT** authorised to administer lidocaine 2% to paediatric patients.