



Clinical Practice Guidelines: Medical/Diabetic emergency: Hyperglycaemia

Policy code	CPG_ME_DHE_0120
Date	January, 2020
Purpose	To ensure consistent management of patients with hyperglycaemia.
Scope	Applies to Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless stated otherwise.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	January, 2023
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by QAS paramedics when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2020.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Diabetic emergency: Hyperglycaemia

January, 2020

A normal fasting blood glucose target for an individual without diabetes is 3.9–5.6 mmol/L. In the diabetic patient hyperglycaemia can present as Diabetic Ketoacidosis (DKA), Hyperosmolar Hyperglycaemic Syndrome (HHS) or be asymptomatic.

DKA and HHS are predominantly caused by:^[1]

- Acute illness (e.g. infection, ACS, CVA)
- Non-compliance with medication.

DKA is a life-threatening complication usually seen in patients with Type 1

Diabetes Mellitus that is characterised by:

- Hyperglycaemia
- Ketosis
- Metabolic acidosis.

It is caused by an **absolute** insulin deficiency, precipitating a number of physiological changes:

- High BGL increases blood osmolarity drawing water out of cells resulting in cellular dehydration.
- High BGL in the kidney filtrate results in osmotic diuresis and polyuria leading to severe dehydration & hypovolaemia. (fluid deficits typically range from 5–8 litres)^[3]
- Alternative fuel sources, including fatty acids, are used, producing organic acids known as ketones. Accumulation of these result in a metabolic acidosis.^[3]
- Dehydration leading to polydipsia.
- Loss of potassium from the body.

HHS is a life-threatening complication of Type 2 Diabetes Mellitus (T2 DM) that is characterised by:

- Hyperglycaemia
- Hyperosmolarity
- Severe dehydration.

NOTE: HHS has been known by numerous names, most notably Hyperosmolar Hyperglycaemic Nonketotic Syndrome (HHNS) and Hyperglycaemic Hyperosmolar Nonketotic Coma (HONK), however HHS is more widely accepted now, as coma is not a prerequisite and patients may present with some degree of ketosis.

HHS is caused by a **relative** insulin deficiency, whereby there is sufficient insulin to limit ketone production thus preventing metabolic acidosis. It most commonly presents in patients > 60 and may be the primary presentation of T2 DM.^[4,5,6] Grossly elevated BGLs still initiate the triad of polyuria, polydipsia and polyphagia, and fluid deficits typically range from 8–10 litres.^[4] HHS has a greater rate of mortality due to the severity of underlying illness, typically sepsis.

Clinical features



The clinical features of DKA and HHS are similar:

Neurological:

- Lethargy
- ALOC
- Seizure
- Coma.

Cardiovascular:

- Signs of hypovolaemia (hypotension, tachycardia)
- Pale, cool or clammy or
- Flushed, hot if febrile.

The exceptions are:

- BGL: DKA (> 10 mmol/L) AND/OR HHS (> 40 mmol/L)
- Kussmaul respiration is due to the severe metabolic acidosis. This is usually not seen in HHS.



Additional information

Mandatory transport criteria

If the patient has ANY of the following, they must be transported to hospital.

- Newly diagnosed diabetes
- Suspected cause of hyperglycaemia is due to the patient being affected by an illness that requires further investigation
- Inability to keep oral fluids down
- Moderate to severe dehydration
- Ketones > 1.5 mmol/L
- Pregnant

Note – Although treatment for severe dehydration may be required, correcting fluid deficits too quickly can cause cerebral oedema – especially in children.^[7,8]

Risk Assessment



- Not applicable

