



Clinical Practice Procedures: Cardiac/Modified Valsalva manoeuvre

| | |
|-----------------------------|---|
| Policy code | CPP_CA_MVM_1017 |
| Date | October, 2017 |
| Purpose | To ensure a consistent procedural approach to modified valsalva manoeuvre. |
| Scope | Applies to Queensland Ambulance Service (QAS) clinical staff. |
| Health care setting | Pre-hospital assessment and treatment. |
| Population | Applies to all ages unless stated otherwise. |
| Source of funding | Internal – 100% |
| Author | Clinical Quality & Patient Safety Unit, QAS |
| Review date | October, 2020 |
| Information security | UNCLASSIFIED – Queensland Government Information Security Classification Framework. |
| URL | https://ambulance.qld.gov.au/clinical.html |

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by QAS paramedics when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2020.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Modified Valsalva manoeuvre

October, 2017

The Valsalva manoeuvre should be the first line of treatment for the management of narrow complex supraventricular tachycardia (SVT) in the haemodynamically stable patient.^[1]

The modified Valsalva manoeuvre with supine repositioning and legs elevated has been shown to be significantly more likely to restore sinus rhythm than the standard Valsalva manoeuvre.^[2]

Given the greater likelihood of success, the modified Valsalva manoeuvre should be performed over standard Valsalva, whenever possible.

Although there are other vagal stimulation methods available, the Valsalva manoeuvre is the safest and most effective technique for terminating SVT of unknown mechanism^[1,3] and is the only QAS approved vagal stimulation method.

Valsalva manoeuvre phases of blood pressure responses

The sinusoidal blood pressure response in normal subjects is divided into four phases:

1. An increase in intrathoracic pressure occurs at the onset of straining. The resulting compressive effect on the aorta increases systolic blood pressure by ≥ 15 mmHg for approximately five seconds.
2. Due to a decrease of venous return, there is a return of systolic blood pressure back to baseline for the remainder of the straining period and an increase in systemic vascular resistance.
3. After the strain is released, the acute decrease in intrathoracic pressure causes an abrupt fall in systolic blood pressure below baseline, and a compensatory increase in heart rate.
4. An increase in venous return causes cardiac output to rise and a corresponding sudden increase in blood pressure that may trigger a reflex bradycardia.

Patients taking beta-blockers typically show a blunted blood pressure response to the Valsalva manoeuvre.

Valsalva manoeuvre heart rate responses for patients with SVT

For a patient with paroxysmal SVT in whom the Valsalva manoeuvre is performed, the potential heart rate and rhythm response outcomes include slowing of sinoatrial nodal activity, block at the atrioventricular node unmasking atrial activity, termination of the SVT, or no response.

Indications

- Haemodynamically stable Supraventricular tachycardia (SVT)

Contraindications^[2]


- Requirement for immediate cardioversion
- Hypotension (SBP < 90 mmHg)
- Atrial fibrillation/flutter
- Aortic stenosis
- Recent myocardial infarction (within 3 months)
- Glaucoma
- Retinopathy
- Third trimester of pregnancy

Complications

- Syncope
- Prolonged hypotensive state

Procedure – Modified Valsalva manoeuvre

1. Obtain a baseline 12-Lead ECG.
2. Explain the procedure to the patient.
3. Position the patient in a semi-recumbent position.



4. Instruct the patient to perform a forced expiration into a sterile 10 mL syringe^[2] for 15 seconds.

6. Reposition the patient to a semi-recumbent position for 45 seconds.
7. Repeat 12-Lead ECG.
8. Confirm the modified Valsalva has been successful. If the patient's SVT has failed to revert, consider repeating the procedure to a maximum of 3 attempts.

Additional information

- If repeated modified Valsalva attempts are required, ensure the patient has returned to a hemodynamically stable SVT presentation prior to repeating.
- The modified Valsalva manoeuvre is the only vagal stimulation technique approved for QAS paramedics.

5. At the end of forced expiration remove the syringe and lay the patient supine with legs raised straight to 45° for 15 seconds.

