



Drug Therapy Protocols: Adrenaline (epinephrine)

Policy code	DTP_ADR_0420
Date	April, 2020
Purpose	To ensure a consistent procedural approach to adrenaline (epinephrine) administration.
Scope	Applies to Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless stated otherwise.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	April, 2023
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.


All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by QAS paramedics when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2020.



This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Adrenaline (epinephrine)

April, 2020

Drug class

Sympathomimetic

Pharmacology

Adrenaline (epinephrine) is a naturally occurring catecholamine which primarily acts on Alpha (α) and Beta (β) adrenergic receptors. The actions of these receptors cause an increase in heart rate (β_1), increase in the force of myocardial contraction (β_1), increase in the irritability of the ventricles (β_1), bronchodilation (β_2) and peripheral vasoconstriction (α_1).^[1-3]

Metabolism

The majority of circulating adrenaline (epinephrine) is metabolised by sympathetic nerve endings. It is subject to the process of mitochondrial enzymatic breakdown by monoamine oxidase at the synaptic level.

Indications

- **Cardiac arrest**
- **Anaphylaxis OR severe allergic reaction**
- **Severe life-threatening bronchospasm**
OR **silent chest** (patients must only be able to speak in single words AND/OR have haemodynamic compromise AND/OR an ALOC)
- **Shock unresponsive to adequate fluid resuscitation** (excluding haemorrhagic, obstructive and anaphylactic)
- **Bradycardia with poor perfusion** (unresponsive to atropine AND/OR transcutaneous pacing)
- **Croup** (moderate to severe)

Contraindications

- Nil

Precautions

- Hypertension
- Hypovolaemic shock
- Concurrent MAOI therapy

Side effects

- Anxiety
- Hypertension
- Palpitations/tachyarrhythmias
- Pupil dilation
- Tremor

Presentation

- Ampoule, 1 mg/1 mL (1:1,000) *adrenaline (epinephrine)*
- Ampoule, 1 mg/10 mL (1:10,000) *adrenaline (epinephrine)*
- Prefilled syringe EpiPen® Auto-injector, 300 microg *adrenaline (epinephrine)*
- Prefilled syringe EpiPen® Jr Auto-injector, 150 microg *adrenaline (epinephrine)*

Onset	Duration	Half-life
30 seconds (IV) 60 seconds (IM)	5–10 minutes	2 minutes

Schedule

- 1 mg/1 mL (1:1,000), S3 (therapeutic poison)
- 1 mg/10 mL (1:10,000), S3 (therapeutic poison)
- 300 microg EpiPen® Auto-injector, S3 (therapeutic poison)
- 150 microg EpiPen® Jr Auto-injector, S3 (therapeutic poison)

Routes of administration

Nebuliser (NEB)	ACP2 CCP
Intramuscular injection (IM)	FR EPTO AT P ACP1 ACP2 CCP
Intravenous injection (IV)	ACP2 CCP
Intraosseous injection (IO)	CCP
Intravenous infusion (IV INF)	CCP
Intraosseous infusion (IO INF)	CCP

Special notes

- Ambulance officers must only administer medications for the listed indications and dosing range. Any consideration for treatment outside the listed scope of practice requires mandatory approval via the QAS Clinical Consult and Advice Line.
- 1:1,000 (1 mg/mL) adrenaline (epinephrine) presentation should be used for all nebuliser administration.
- 1:10,000 (100 microg/1 mL) or a 1:100,000 (10 microg/1 mL) adrenaline (epinephrine) preparation should be used for all low dose IM/IV injections. Ensure all syringes are appropriately labelled.
- If possible, all time critical adrenaline (epinephrine) IM injections should be administered in the vastus lateralis (improved absorption).
- Suitably qualified officers should, whenever possible, administer adrenaline infusions through an appropriately placed central venous line.
- Suitably qualified officers should, whenever possible, use invasive pressure monitoring for patients being administered adrenaline (epinephrine) infusions.
- Adrenaline (epinephrine) infusions must be administered through a dedicated line.
- Patients on adrenaline (epinephrine) infusions without continuous IBP monitoring **must** have their NIBP measured regularly (every 5 mins at a minimum).
- All cannulae with adrenaline (epinephrine) infusions should be as proximal as possible, be freely flowing, and be watched closely for extravasation.
- NIBP cuffs **must not be** placed on limbs with infusions to ensure flow is not obstructed.
- All cannulae and IV lines must be flushed thoroughly with sodium chloride 0.9% following each medication administration.

Adult dosages

Cardiac arrest		
ACP2 CCP	IV	1 mg Repeated at 3 – 5 minute intervals. No maximum dose.
CCP	IO	1 mg Repeated at 3 – 5 minute intervals. No maximum dose.

Anaphylaxis OR severe allergic reaction		
FR EPTO AT P	IM	EpiPen® Auto-injector (300 microg) Single dose only.
ACP1 ACP2 CCP	IM	500 microg (300 microg if known pregnancy) Repeated at 5 minute intervals. No maximum dose.
ACP2 CCP	NEB	5 mg Single dose only. May be administered for upper airway obstruction thought to be allergic in origin – IM adrenaline (epinephrine) must first be administered.
CCP	IV/IO INF	May be administered for refractory anaphylaxis unresponsive to 3 x IM adrenaline (epinephrine) injections and adequate fluid administration. 20 – 50 microg bolus (IV/IO) Immediately followed by an infusion commencing at 10 microg/minute (10 mL/hr) – titrate accordingly to indication and patient’s physiological response to treatment. Maximum infusion rate 50 microg/min (50 mL/hr). <i>Infusion preparation: Mix 3 mg (3 mL) of 1:1000 adrenaline (epinephrine) with 47 mL of sodium chloride 0.9% to achieve a final concentration of 60 microg/mL. Ensure the syringe is appropriately labelled. Administer the infusion via the Perfusor® Space Medication Library (Adrenaline-Adult (shock)).</i>

Severe life-threatening bronchospasm OR silent chest
(patients must only be able to speak in single words AND/OR have haemodynamic compromise AND/OR an ALOC)

ACP1 ACP2 CCP	IM	500 microg (300 microg if known pregnancy) Repeated at 5 minute intervals. No maximum dose.
CCP	IV/IO INF	May be administered for refractory severe life-threatening bronchospasm or silent chest unresponsive to 3 x IM adrenaline (epinephrine) injections and adequate fluid administration. 20 – 50 microg bolus (IV/IO) Immediately followed by an infusion commencing at 10 microg/minute (10 mL/hr) – titrate accordingly to indication and patient’s physiological response to treatment. Maximum infusion rate 50 microg/min (50 mL/hr). <i>Infusion preparation: Mix 3 mg (3 mL) of 1:1000 adrenaline (epinephrine) with 47 mL of sodium chloride 0.9% to achieve a final concentration of 60 microg/mL. Ensure the syringe is appropriately labelled. Administer the infusion via the Perfusor® Space Medication Library (Adrenaline-Adult (shock)).</i>

Shock unresponsive to adequate fluid resuscitation
(excluding haemorrhagic, obstructive and anaphylactic)

CCP	IV/IO INF	20 – 50 microg bolus (IV/IO) Immediately followed by an infusion commencing at 10 microg/minute (10 mL/hr) – titrate accordingly to indication and patient’s physiological response to treatment. Maximum infusion rate 50 microg/min (50 mL/hr). <i>Infusion preparation: Mix 3 mg (3 mL) of 1:1000 adrenaline (epinephrine) with 47 mL of sodium chloride 0.9% to achieve a final concentration of 60 microg/mL. Ensure the syringe is appropriately labelled. Administer the infusion via the Perfusor® Space Medication Library (Adrenaline-Adult (shock)).</i>
-----	--------------	---

Adult dosages (cont.)

Bradycardia with poor perfusion

(unresponsive to atropine AND/OR transcutaneous pacing)

CCP	IV/IO	20 – 50 microg Repeated at 1 minute intervals. No maximum dose.
-----	-------	---

Paediatric dosages

Cardiac arrest

ACP2 CCP	IV	10 kg or more (≥ 1 year) – 10 microg/kg Repeated at 3 – 5 minute intervals. No maximum dose. Less than 10 kg (excluding newly born) – 100 microg Repeated at 3 – 5 minute intervals. No maximum dose. Newly born – 50 microg Repeated at 3 – 5 minute intervals. No maximum dose.
CCP	IO	10 kg or more (≥ 1 year) – 10 microg/kg Repeated at 3 – 5 minute intervals. No maximum dose. Less than 10 kg (excluding newly born) – 100 microg Repeated at 3 – 5 minute intervals. No maximum dose. Newly born – 50 microg Repeated at 3 – 5 minute intervals. No maximum dose.

Paediatric dosages (cont.)

Anaphylaxis OR severe allergic reaction

FR EPTO AT P	IM	6 years or older – EpiPen® Auto-injector (300 microg). Single dose only. 1 year–less than 6 years – EpiPen® Jr Auto-injector (150 microg)
ACP1 ACP2 CCP	IM	6 years or older – 300 microg Repeated at 5 minute intervals. No maximum dose. 1 year – less than 6 years – 150 microg Repeated at 5 minute intervals. No maximum dose. 6 months – less than 1 year – 100 microg Repeated at 5 minute intervals. No maximum dose. Less than 6 months – 50 microg Repeated at 5 minute intervals. No maximum dose.
ACP2 CCP	NEB	5 mg Single dose only. May be administered for upper airway obstruction thought to be allergic in origin – IM adrenaline (epinephrine) must first be administered.
CCP	IV/IO INF	May be administered for refractory anaphylaxis unresponsive to 3 x IM adrenaline (epinephrine) injections and adequate fluid administration. 6 years or older – 1 microg/kg bolus (IV/IO) Immediately followed by an infusion commencing at 0.2 microg/kg/min (0.2 mL/hr) – titrate accordingly to indication and patient’s physiological response to treatment. Maximum infusion rate 0.5 microg/kg/min Less than 6 years – QAS Clinical Consultation and Advice Line consultation and approval required in all situations. Infusion preparation: Mix 3 mg (3 mL) of 1:1000 adrenaline (epinephrine) with 47 mL of sodium chloride 0.9% to achieve a final concentration of 60 microg/mL. Ensure the syringe is appropriately labelled. Administer the infusion via the Perfusor® Space Medication Library (Adrenaline-Paed (Shock)).

Paediatric dosages (cont.)

Severe life-threatening bronchospasm OR silent chest

(patients must only be able to speak in single words AND/OR have haemodynamic compromise AND/OR an ALOC)

ACP1 ACP2 CCP	IM	<p>6 years or older – 300 microg Repeated at 5 minute intervals. No maximum dose.</p> <p>1 year – less than 6 years – 150 microg Repeated at 5 minute intervals. No maximum dose.</p> <p>6 months – less than 1 year – 100 microg Repeated at 5 minute intervals. No maximum dose.</p> <p>Less than 6 months – 50 microg Repeated at 5 minute intervals. No maximum dose.</p>
CCP	IV/IO INF	<p>May be administered for refractory severe life-threatening bronchospasm or silent chest unresponsive to 3 x IM adrenaline (epinephrine) injections and adequate fluid administration.</p> <p>6 years or older – 1 microg/kg bolus (IV/IO) Immediately followed by an infusion commencing at 0.2 microg/kg/min (0.2 mL/hr) – titrate accordingly to indication and patient’s physiological response to treatment. Maximum infusion rate 0.5 microg/kg/min</p> <p>Less than 6 years – QAS Clinical Consultation and Advice Line consultation and approval required in all situations.</p> <p><i>Infusion preparation: Mix 3 mg (3 mL) of 1:1000 adrenaline (epinephrine) with 47 mL of sodium chloride 0.9% to achieve a final concentration of 60 microg/mL. Ensure the syringe is appropriately labelled. Administer the infusion via the Perfusor® Space Medication Library (Adrenaline-Paed (shock)).</i></p>

Croup (moderate to severe)

ACP2 CCP	NEB	5 mg Single dose only.
-------------	-----	---------------------------

Shock unresponsive to adequate fluid resuscitation

(excluding haemorrhagic, obstructive and anaphylactic)

CCP	IV/IO INF	QAS Clinical Consultation and Advice Line consultation and approval required in all situations.
-----	--------------	---

Bradycardia

(unresponsive to atropine AND/OR transcutaneous pacing)

CCP	IV/IO	QAS Clinical Consultation and Advice Line consultation and approval required in all situations.
-----	-------	---