



Drug Therapy Protocols: Naloxone

Policy code	DTP_NAL_0519
Date	May, 2019
Purpose	To ensure a consistent procedural approach to naloxone administration.
Scope	Applies to all Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless specifically mentioned.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	May, 2022
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While effort has been made to contact all copyright owners this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by QAS paramedics when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2019.



This work is licensed under the **Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License**

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au



Drug class

Opioid antagonist

Pharmacology

Naloxone is an opioid antagonist that prevents or reverses the effects of opioids including respiratory depression, sedation and hypotension. Naloxone antagonises the opioid effects by competing for the same receptor sites.^[1-3]

Metabolism

Hepatic.^[1]

Indications

- **Respiratory depression** (secondary to the administration of narcotic drugs)

Contraindications

- Allergy and/or Adverse Drug Reaction
- The newly born patient

Precautions

- Use with caution on patients with pre-existing cardiac disease

Side effects

- Narcotic reversal can cause combativeness, vomiting, sweating, tachycardia and hypertension
- May produce acute withdrawal convulsions in the chronic narcotic user
- Pulmonary oedema

Presentation

- Ampoule, 400 microg/mL *naloxone hydrochloride dihydrate*

Onset

3–5 minutes (IM)
/1–3 minutes (IV)

Duration

≈ 60 minutes

Half-life

60 minutes


Schedule

- S3 (Pharmacist Only).

Routes of administration

Intramuscular injection (IM)

Intravenous injection (IV)



Special notes ^[4,5,6]

- Ambulance officers must only administer medications for the listed indications and dosing range. Any consideration for treatment outside the listed scope of practice requires mandatory approval via the QAS Clinical Consult and Advice Line.
- Naloxone should only be administered following adequate patient oxygenation and ventilation.
- Naloxone should be administered cautiously to patients who are known or suspected to be physically dependent on narcotics.
- In the vast majority of cases, naloxone should not be required and the patient will need only supportive therapy followed by transport to a medical facility.
- The duration of the narcotic may exceed that of naloxone and renarcotisation is always a possibility.
- Administration of naloxone in the pre-hospital environment may unmask potentially unwanted side effects in the setting of poly pharmacy overdose.
- Naloxone should not be administered to the newly born, even in the setting of suspected or identified opiate exposure/overdose, as administration of naloxone may lead to acute withdrawal and seizures.

Adult dosages

Respiratory depression (secondary to the administration of narcotic drugs)

ACP1 ACP2 CCP	IM	1.6 mg Single dose only.
CCP	IV	50 microg Repeated PRN to facilitate airway management. No maximum dose.

Paediatric dosages

Respiratory depression (secondary to the administration of narcotic drugs)

Single dose of naloxone to be administered using the following scale:

Weight	Dose	Volume
< 5 kg	100 microg	0.25 mL
5 – < 10 kg	200 microg	0.5 mL
10 – < 15 kg	300 microg	0.75 mL
15 – < 20 kg	400 microg	1 mL
20 – < 25 kg	500 microg	1.25 mL
25 – < 30 kg	600 microg	1.5 mL
30–35kg	700 microg	1.75 mL
> 35 kg	800 microg	2 mL

ACP2
CCP

IM

20 microg/kg
Single dose only, not to exceed 800 microg.